



CrossMark  
click for updates

## Introduction

**Cite this article:** Sammonds P, Montagnat M, Bons P, Schneebeli M. 2017 Ice microstructures and microdynamics. *Phil. Trans. R. Soc. A* **375**: 20160438.  
<http://dx.doi.org/10.1098/rsta.2016.0438>

Accepted: 7 November 2016

One contribution of 11 to a theme issue  
'Microdynamics of ice'.

### Subject Areas:

glaciology

### Keywords:

ice microstructures, ice cores, microdynamics, recrystallization, scaling

### Author for correspondence:

Peter Sammonds

e-mail: [p.sammonds@ucl.ac.uk](mailto:p.sammonds@ucl.ac.uk)

# Ice microstructures and microdynamics

Peter Sammonds<sup>1</sup>, Maurine Montagnat<sup>2</sup>, Paul Bons<sup>3</sup>  
and Martin Schneebeli<sup>4</sup>

<sup>1</sup>Rock and Ice Physics Laboratory, Department of Earth Sciences, University College London, London WC1E 6BT, UK

<sup>2</sup>Laboratoire de Glaciologie et Géophysique de l'Environnement, CNRS, St Martin d'Hères cedex, France

<sup>3</sup>Department of Geosciences, Eberhard Karls University Tübingen, Wilhelmstrasse 56, Tübingen 72074, Germany

<sup>4</sup>WSL Institute for Snow and Avalanche Research SLF, Davos 7260, Switzerland

 PS, 0000-0003-3355-6761

## 1. Introduction

Ice occurs as polycrystalline aggregates in which the bulk behaviour is the result of the behaviour of the ensemble of individual grains, and is therefore dependent on the microstructure, that is to say, the whole arrangement of grains (size, shape, and shape and lattice orientations), their internal substructure (dislocation density and subgrains), impurities and second phases (bubbles in firn, clathrates and dissolved impurities). Knowledge of ice microstructures is central to our understanding of ice microdynamics. It is also important for our understanding of ice in relation to other materials, such as metals and rocks, and of the controls microstructures exert on macro-scale deformation processes in glaciers, polar ice caps, the sea ice cover and icy planets. The microstructure is a constantly changing and evolving, dynamic entity. Two important processes that change the microstructure are grain growth (driven by surface-energy reduction) and (discontinuous) recrystallization (driven by strain energy); others include the plastic deformation and fracture of asperities of ice. These are the foci of current research. This theme issue covers glacier ice,

polar ice, sea ice and planetary ices, with cross-cutting themes of ice microstructures and recrystallization, ice-core studies, modelling of constitutive laws, planetary ices and scaling. But the underlying relationship examined is that between microstructures, fabrics and asperities, and deformation. Although microstructural dynamics impacts on almost every aspect of ice behaviour, the topics covered in this theme issue are currently at the forefront of international research efforts.

## 2. Recrystallization

Recrystallization is of particular interest in materials science as recrystallization dramatically changes the properties of a material, notably the mechanical properties such as strength, ductility and anisotropy. Dynamic recrystallization processes in ice are similar to those observed in rocks and metals [1]. Chauve *et al.* [2] describe detailed experiments and microstructural studies investigating nucleation processes during dynamic recrystallization of ice using cryogenic electronic backscattered diffraction, while Llorens *et al.* [3] describe the dynamic recrystallization during deformation of polycrystalline ice through insights derived from numerical simulations. The physical processes involved in recrystallization are nucleation of recrystallized grains by local dislocation rearrangements and growth of these grains by grain boundary migration with concurrent consumption of the deformation-induced defects by the moving grain boundaries. Since nucleation is triggered by the heterogeneity of the deformed structure and, therefore, at sites which are not typical for the homogeneously deformed material, recrystallization kinetics and microstructure evolution are extremely difficult to model or predict. This is the reason that recrystallization is often referred to as the last unsolved problem in physical metallurgy and that further research is still necessary although the basic phenomenon has been treated in the scientific literature for more than 120 years. Solving this problem is as relevant for ice as it is for metals. The flow of glaciers and polar ice sheets is controlled by the highly anisotropic rheology of ice that is close to its melting point. During deformation within an ice sheet, recrystallization processes occur that modify the ice texture and microstructure and, therefore, affect the flow behaviour. Dynamic recrystallization controls ice microstructures and rheology under different boundary conditions that range from pure shear flattening at the top to simple shear near the base of the sheets. Llorens *et al.* [3] have investigated these processes through simulations of viscoplastic deformation in combination with dynamic recrystallization in ice.

Recrystallization is intentionally used in engineering to optimize materials properties of steels and non-ferrous alloys [4]. The interplay of forming strain, temperature and time generates a wide spectrum of different microstructures that are associated with different mechanical properties. Metals, rocks and ice, all polycrystalline aggregates, show remarkable similarities in their recrystallization behaviour. In this respect, hot rolling of a steel sheet, the formation of a mountain belt and the flow of a glacier are, therefore, highly comparable processes, despite the differences in length and time scale, and the material-specific properties of the individual polycrystals. This has an echo in the study described in this volume of recrystallization and damage of ice caused by friction of runners against ice in winter sports by Seymour-Pierce *et al.* [5]. Ice-specific complications arise, for example, in sea ice, where a second, saline, phase may interact with the polycrystal and significantly modify its microstructure, and hence mechanical and physical behaviour. An example is shown in the study by Marchenko & Lishman [6] on the influence of closed brine packets and permeable brine channels on the thermo-elastic properties of saline ice.

## 3. Ice-core studies

Ice sheets allow us to extract hundreds of thousands of years of climate record. However, complex deformation processes occur and still hamper accurate dating of events. This uncertainty in the

ice record can be drastic, particularly for the deepest and oldest ice. For the reconstruction of our planet's climate history, it is imperative that the mechanisms and processes, such as flow and recrystallization, that modify this record are better understood and constrained. The variation of microstructures along ice cores also provides a unique image of the flow history of ice caps and glaciers. For example, long-favoured models on the stratification of ice caps are currently being challenged and improved, partly owing to new developments in the analysis of ice-core microstructures (e.g. microstructure mapping) and numerical modelling techniques (e.g. micromechanical modelling). Weikusat *et al.* [7] describe the physical analysis of an Antarctic ice core drilled at Kohnen Station, Dronning Maud Land, Antarctica, as a move towards an integration of the micro- and macrodynamics of polar ice.

## 4. Constitutive laws for ice

Ice-sheet modelling has nearly always assumed a constitutive response based on an incompressible, nonlinearly viscous law with a temperature-dependent rate factor, and used a reduced model (shallow ice approximation) for the momentum and energy balances. Furthermore, the viscous response is commonly taken to be a power law, based on Glen's 1955 data. Although first-order approximations for constitutive laws have significantly advanced our knowledge of the behaviour of ice, more accurate descriptions of the behaviour of ice are needed, for example, to develop predictive models for the future of our planet's ice caps, sea level and climate. Again, the presence of impurities or seawater in pore spaces will have a major impact on flow behaviour. Middleton *et al.* [8] describe the influence of a particulate mixture of fluorite in ice. Ice-rock mixtures are found in a range of natural terrestrial and planetary environments.

While the bulk flow properties of ice are important, so too are the friction properties of ice in the Earth dynamics, for instance in glacier and ice stream flow over bedrock and sediments, in the dynamics of the Arctic Ocean, and also of course in winter sports! Three papers in this theme issue address friction. McCarthy *et al.* [9] use an empirical approach of rate and state friction, with parameters derived from laboratory ice-rock friction studies to arrive at a constitutive relation applicable to glacier motion over bedrock. Sammonds *et al.* [10] test a microphysics- and micromechanics-based constitutive friction law against intermediate-scale experiments on saline ice floes conducted in an environmental basin. Seymour-Pierce *et al.* [5] consider an energy balance approach when modifying Amontons' law to model the sliding of runners in the skeleton winter sport.

Close collaboration is needed between microstructure researchers, engineers, planetary scientists and geophysicists to improve constitutive laws. Whatever constitutive law is adopted, the numerical problems of solving three-dimensional unsteady flow equations (thermo-mechanically coupled momentum and energy balances on an unknown evolving domain), real ice-sheet modelling, will be the next formidable challenge.

## 5. Planetary ices

The exploration of the Martian surface by the rover Spirit, the fly past of Jupiter's icy moon Europa by Galileo and Voyager reaching the distant world of Neptune's moon Triton and beyond (e.g. the Cassini-Huygens mission to Titan) have not only thrilled the public but have produced images of such high resolution that scientists are beginning to unravel the tectonics of these planets with ever greater confidence. Throughout the Solar System ices exist. However, their compositions vary widely. For instance, hydrated magnesium sulfate minerals, such as kieserite ( $\text{MgSO}_4 \cdot \text{H}_2\text{O}$ ), hexahydrate ( $\text{MgSO}_4 \cdot 6\text{H}_2\text{O}$ ), epsomite ( $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ ) and meridianiite ( $\text{MgSO}_4 \cdot 11\text{H}_2\text{O}$ ), are likely to be important planetary materials, as are clathrate hydrates. On the icy moons, these hydrated phases have been suggested as possible non-water phases at the surface, and may affect the possibility and rate of solid-state convection within the interiors. On Mars, meridianiite may be the most abundant hydrate mineral on the planet, and is important in studies of the present Martian water budget. While the fundamental laws of mechanics will apply whatever

the situation, what differs are the driving forces and the response of planetary materials because of the different environmental conditions to which they are subject. Grain-size-dependent flow in Ice I has been suggested as playing an important role on the Earth [11]. However, it has now been shown that a high-pressure polymorph of ice, Ice II, an important planetary-forming mineral, also deforms by grain-size sensitive creep at low stresses, while the effect of particulate matter on behaviour at high homologous temperatures has been shown to be important during deformation [12]. The effect of the increase in strength due to increasing rock particle content [8] will impact models of planetary evolution, while the analyses of deformation fabrics of candidate planetary ices at relevant pressure–temperature conditions are going to be key to understanding their rheology.

## 6. Scaling

The final topic of this theme issue is the state of the art in the understanding of microdynamics of ice in relation to the controls microstructures exert on large-scale deformation processes in glaciers, polar ice caps, the sea ice cover and the icy planets. Weiss & Dansereau [13] analyse the linking across scales for sea ice mechanics. As they discuss, this question is not new [14], but has re-emerged because of the analysis of *in situ* and remote sensing data of the Arctic sea ice cover, and it is importance in understanding the dynamical response of the Arctic to global warming. The analysis of heterogeneous media is generally handled through homogenization procedures. But issues can arise with this approach [13]. Scaling is also addressed by Sammonds *et al.* [10] and Weikusat *et al.* [7] and indirectly by many of the contributors to this theme issue. Understanding the key underlying relations between microstructures and fabrics and deformation is important for improved large-scale models of glaciers, polar ice, the Arctic ice cover and the icy planets.

**Competing interests.** We declare we have no competing interests.

**Funding.** No funding has been received for this article.

## References

1. Wilson CJL, Peternell M, Piazoloc S, Luzind V. 2014 Microstructure and fabric development in ice: lessons learned from in situ experiments and implications for understanding rock evolution. *J. Struct. Geol.* **61**, 50–77. (doi:10.1016/j.jsg.2013.05.006)
2. Chauve T, Montagnat M, Barou F, Hidas K, Tommasi A, Mainprice D. 2017 Investigation of nucleation processes during dynamic recrystallization of ice using cryo-EBSD. *Phil. Trans. R. Soc. A* **375**, 20150345. (doi:10.1098/rsta.2015.0345)
3. Llorens M-G, Griera A, Steinbach F, Bons PD, Gomez-Rivas E, Jansen D, Roessiger J, Lebensohn RA, Weikusat I. 2017 Dynamic recrystallization during deformation of polycrystalline ice: insights from numerical simulations. *Phil. Trans. R. Soc. A* **375**, 20150346. (doi:10.1098/rsta.2015.0346)
4. Ashby M, Shercliff H, Cebob D. 2014 *Materials: engineering, science, processing and design*. Amsterdam, The Netherlands: Elsevier.
5. Seymour-Pierce A, Lishman B, Sammonds P. 2017 Recrystallization and damage of ice in winter sports. *Phil. Trans. R. Soc. A* **375**, 20150353. (doi:10.1098/rsta.2015.0353)
6. Marchenko A, Lishman B. 2017 The influence of closed brine packets and permeable brine channels on the thermo-elastic properties of saline ice. *Phil. Trans. R. Soc. A* **375**, 20150351. (doi:10.1098/rsta.2015.0351)
7. Weikusat I *et al.* 2017 Physical analysis of an Antarctic ice core—towards an integration of micro- and macrodynamics of polar ice. *Phil. Trans. R. Soc. A* **375**, 20150347. (doi:10.1098/rsta.2015.0347)
8. Middleton CA, Grindrod PM, Sammonds PR. 2017 The effect of rock particles and D<sub>2</sub>O replacement on the flow behaviour of ice. *Phil. Trans. R. Soc. A* **375**, 20150349. (doi:10.1098/rsta.2015.0349)

9. McCarthy C, Savage H, Nettles M. 2017 Temperature dependence of ice-on-rock friction at realistic glacier conditions. *Phil. Trans. R. Soc. A* **375**, 20150348. (doi:10.1098/rsta.2015.0348)
10. Sammonds PR, Hatton DC, Feltham DL. 2017 Micromechanics of sea ice frictional slip from test basin scale experiments. *Phil. Trans. R. Soc. A* **375**, 20150354. (doi:10.1098/rsta.2015.0354)
11. Goldsby DL, Kohlstedt DL. 1997 Grain boundary sliding in fine-grained Ice I. *Scr. Mater.* **37**, 1399–1406. (doi:10.1016/S1359-6462(97)00246-7)
12. Arenson LU, Springman SM, Segal DC. 2007 The rheology of frozen soils. *Appl. Rheol.* **17**, 12147-1–12147-14. (doi:10.3933/ApplRheol-17-12147)
13. Weiss J, Dansereau V. 2017 Linking scales in sea ice mechanics. *Phil. Trans. R. Soc. A* **375**, 20150352. (doi:10.1098/rsta.2015.0352)
14. Overland JE, Walter BA, Curtin TB, Turet P. 1995 Hierarchy and sea ice mechanics: a case study from the Beaufort Sea. *J. Geophys. Res.* **100**, 4559–4571. (doi:10.1029/94JC02502)